

# **Lattice-Automaton Modelling of Bioturbation and Benthic Activity**

Bernard P. Boudreau  
NERC Challenger Division for Sea Floor Processes  
Southampton Oceanography Centre, University of Southampton  
European Way, Southampton SO14 3ZH United Kingdom

and currently:

Department of Oceanography, Dalhousie University  
Halifax, Nova Scotia B3H 4J1, Canada  
phone: (902) 494-8895 fax: (902) 494-3877 e-mail: [bpboudr@is.dal.ca](mailto:bpboudr@is.dal.ca)

Grant Number: N000140110250

## **LONG-TERM GOALS**

My long term goal is a quantitative and mechanistic understanding of the relationship between infaunal ecology, their actions and their rates and the consequent modification of sediments, including the creation and destruction of heterogeneities and the modes and rates of sediment component mixing.

## **OBJECTIVES**

The development of a model/computer code that embraces the discrete nature of sediments and organisms, rather than averaging it away, and that utilizes biologically relevant parameters, such as animal sizes, population density, feeding and locomotion rates, and probabilities for observed behaviour(s), to drive the model and produce predictions about sediment composition and fabric.

## **APPROACH**

My approach is the direct modelling of organism-sediment interactions via a new type of model. Biologically active sediment is represented on a computer as a regular lattice of quasi-particles with individually assigned chemical, biological or physical properties. Model benthic organisms are introduced in the form of automaton, i.e. programmable entities, that are capable of moving through the particle lattice by displacing or ingesting-defecating particles. Each automaton obeys a set of rules, both deterministic and stochastic, designed to mimic real organism behavior, and different types of organisms have different sets of rules.

## **WORK COMPLETED**


The computer code of this model for small deposit feeders, which carries the acronym LABS, is complete and extensive study of the effect of these organisms on tracer distributions has been completed and is now in press. (This constitutes the validation phase of the project.)

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>30 SEP 2001</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2001 to 00-00-2001</b>	
4. TITLE AND SUBTITLE <b>Lattice-Automaton Modelling of Bioturbation and Benthic Activity</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>NERC Challenger Division for Sea Floor Processes,,Southampton Oceanography Centre, University of Southampton,,European Way, Southampton SO14 3ZH United Kingdom, ,</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>7</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			


## RESULTS

The results from the model are, so far, of two types: fabric representations and tracer distributions.

*Fabric:* 2D visualizations of state of a sediments particle distribution provide immediate appreciation of the consequences of animal actions on sediment fabric and composition, including the development of biologically-induced heterogeneities, which are observed in real sediments. Figure 1A illustrates an initial random fabric with four worms beginning to process the sediment; Fig. 1B shows the effects of burrowing and packaging after 100 days, i.e., the sediment fabric is now dominated by the biological features.




***Figure 1A. Sediment with an initial random distribution of particles (white) and water (blue), with porosity of 0.8, and distinct sediment-water interface. Four small deposit-feeding worms (yellow with red heads) are beginning to rework the sediment. (Sediment is 50 cmX25 cm, worms are 1-4 cm in length.)***



***Figure 1B. Same sediment after 1 year of reworking. Note burrows and heterogeneities.***


These results (Fig. 1) allow study of both the size and the rate of change of the heterogeneity with time and as a function of animal parameters, such as size and number.

Tracer Distributions: Each particle in the model can be tagged with a selected amount of tracer, and this can be modified with time. Thus, it is possible to simulate the arrival of particle specific tracers, like  $^{210}\text{Pb}$ , that are used to estimate mixing rates in natural sediments. From the model 2D distribution of  $^{210}\text{Pb}$ , it is then possible to create a synthetic 1D  $^{210}\text{Pb}$  profile with depth (Figure 2), and from that profile calculate apparent mixing coefficient values,  $D_B$ , as is done for real sediments.



**Figure 2.** Synthetic  $^{210}\text{Pb}$  profiles obtained by laterally averaging the  $^{210}\text{Pb}$  concentrations in the 2D model (one organism, 0.0cm/yr sedimentation, 1-10 g/d feeding rate, 1-10 cm/d locomotion rate.) The steady state profile at 100 years (pink-violet) can be modelled with a standard diffusion model to obtain  $D_B$  values.

All model runs of LABS are done with literature quoted biological parameter values, without any a priori adjustments. Typical values of  $D_B$  as a function of animal locomotion speed (with and without ingestion-egestion), animal numbers, mouth size, are shown in Figure 3 (see also Boudreau et al., in press, as listed below, for a more extensive discussion of these results). The range of  $D_B$  values is very much that seen in real sediments, which validates the model and illustrates for the first time the link between specific animal activities and  $D_B$ .




**Figure 3. Plots showing the dependence of  $D_B$  on locomotion, number of organisms, mixing depth and organism width, as observed in the small deposit-feeder implementation of the LABS code.**

Conversely to a specific steady state tracer, it is possible to label all particle with a number and simply calculate  $D_B$  from the time-dependent standard-deviation (displacements) of the entire population of particles,  $\overline{\Delta^2}$ ,

$$\overline{\Delta^2} = 4 D_B t$$

Such a function is plotted in Figures 4. The trend in the results is linear for times greater or equal to about 1 year. The calculated  $D_B$  values are in line with the  $^{210}\text{Pb}$  results. At early times, however, the trend is strongly non-linear, indicating that  $D_B$  is then a function of time. This has important implications for some types of tracer studies where an experimenter salts the sediment-water interface

with tagged particles. Such experiment must have a duration of at least 9-12 months if the true  $D_B$  value is to be obtained, and this has not always been the case.



**Figure 4.** Plot of the root-mean-square displacement of all the particles in a 2D simulation as a function of time. Left-hand diagram is for the entire time ranged used, while the right-hand diagram is for early times.

Even at this early stage, the lattice-automaton model is providing significant insight into the working of bioturbation and our standard means of measuring and quantifying these effects.

### IMPACT/APPLICATIONS

The model needs to be expanded to a broader ecology of infaunal functional types before it can be considered to be a realistic simulation of sediments. Nevertheless, the results so far are highly encouraging, and we may soon have a computer-based method of simulating relatively complete benthic ecologies and their effects.

### TRANSITIONS

The LABS has been given to my NICOP partners and will soon be in the public domain via publications listed below.

### RELATED PROJECTS

Larry Mayer and Peter Jumars at the Darling Center are collecting data we can use to extend the model, including transformation of sediments by passage through an animal’s gut.

## **PUBLICATIONS**

Choi, J., François-Carcaillet, F. and Boudreau, B.P. (in press) Lattice-automaton bioturbation simulator (LABS): Implementation for small deposit feeders. *Computers and Geosciences*, October issue.

Boudreau, B.P., Choi, J. and François-Carcaillet, F. (in press) Diffusion in a lattice-automaton model of bioturbation by small deposit feeders. *Journal of Marine Research*, september issue.